



REVIEW ARTICLE

Botany and Ethnopharmacological Potential of Ashwagandha

Arjun Pandian^{1*}, Kaliyaperumal Ashokkumar^{2*}, Sudharshan Sekar³, P. Sivakumar⁴, K.S. Vijai Selvaraj⁴, M. Karthik⁵ and L. Hariprasath⁶

¹Department of Plant Biotechnology, PRIST Deemed University, Vallam, Thanjavur, Tamil Nadu, India

²Cardamom Research Station, Kerala Agricultural University, Pampadumpara, Idukki, Kerala, India

³Department of Biotechnology and Food Technology, University of Johannesburg, Auckland Park, South Africa

⁴Agriculture College and Research Institute, TNAU, Eachangkottai, Thanjavur, Tamil Nadu, India

⁵Department of Agricultural Microbiology, TNAU, Coimbatore, Tamil Nadu India.

⁶Department of Biochemistry, Karpagam University, Coimbatore, Tamil Nadu, India

ABSTRACT

Edited by:

Dr. K.Ashokkumar, Ph.D.,
Cardamom Research Station, KAU,
Pampadumpara, Idukki, Kerala.

Reviewed by:

Dr. Raju Ramasubbu,
Department of Biology,
Gandhigram Rural Institute,
Gandhigram, Dindigul, TN, India

Dr. S. Ramesh, Ph.D.,
Cardiovascular and Mitochondrial
Related Disease Research Center,
Hualien Tzu Chi Hospital,
Buddhist Tzu Chi Medical Foundation,
Hualien 970, Taiwan

Article history:

Received: December 05, 2020

Accepted: December 29, 2020

Published: December 30, 2020

Citation:

Pandian, A., Ashokkumar, K., Sekar, S., ,
Sivakumar, P., Selvaraj, K.S.V., Karthik,
M. & Hariprasath, L. (2020). Botany
and ethnopharmacological potential of
Ashwagandha. *Journal of Current
Opinion in Crop Science*, 1(1), 35-40.

Ashwagandha [*Withania somnifera* (L.) Dunal] roots have been used for Indian traditional medicine applications including for the control of fever, asthma, arthritis, rheumatism, inflammation, tuberculosis, mental diseases, and male sexual disorders. The multipurpose use of Ashwagandha has numerous other beneficial health effects that are relevant in light of pharmaceutical perspectives. This review aims to provide comprehensive evaluation of the botanical description and ethnobotanical uses of ashwagandha. An online survey was accompanied for traditional applications of ashwagandha extracts. Appropriate data were obtained from several electronic scientific databases and additional information was also obtained from other relevant sources like textbooks and local prints and scripts. This review investigates and presents the relevant information on ashwagandha, and its botany, ethnopharmacological uses. Future online survey studies are needed to understand the bioactive mechanism of alkaloids, and steroidal lactones compounds, and their effects upon their or consumption through which we can facilitate the protection of human health from several diseases.

Keywords: Ashwagandha; Botany; Ethnopharmacology;
Traditional medicine

*Corresponding author e-mail address: arjungri@gmail.com (A. Pandian) and biotech.ashok@gmail.com (K. Ashokkumar)

INTRODUCTION

Withania somnifera (L.) Dunal commonly known as Ashwagandha, belongs to family Solanaceae, is an important Ayurvedic medicinal plant, its pharmacological, chemical composition and therapeutic effectiveness has been recognized. Ashwagandha is commonly known as Indian winter cherry and Indian ginseng, in India it's called by different vernacular names like Amukkira (Tamil), Asgandh (Hindi), Tilli (Marathi) and Akshan (Pujabi). For centuries, parts of the plant have been used for treating several diseases by different medical systems, depending on ethnic groups. In traditional Indian medicines, this plant has grouped as Rasayana (lengthening lifespan) in Ayurveda. Daily consumption of powdered root of this plant is assumed to delay the senescence, rejuvenate the muscle and reproductive organs and increase the fertility (Durg et al., 2018). In folk medicine, it's used to treat cough, stress, nervous exhaustion, memory loss and muscular weakness (Sharma and Dash, 1998).

The root extracts of the plant exposed to be efficiently improved cerebral health, brain ageing and locomotive syndrome (Singh et al., 2008). It's also used to cure weakness, epilepsy, memory loss, and neurodegenerative diseases such as spinal cord injury, Alzheimer's, and Parkinson's disease (Kulkarni and Dhir, 2008; Rajasankar et al., 2009). Additionally, ashwagandha root extracts are used or consumed as a dietary supplement across developing the world as well as developed nation like the United States (Rajasankar et al., 2009). Based on the above potential health benefits, ashwagandha had great attention in traditional medicine. Therefore, the aim of this review is to highlight the main phytochemicals and its beneficial effects of ashwagandha root powder and extracts for human wellness.

Botanical description

Ashwagandha (*Withania somnifera* L. Dunal) belongs to the Solanaceae family. The plant grows well in dry and arid soils of subtropical countries of Asia and Africa continents (Atal and Schwarting, 1961). In India, ashwagandha prolifically grows in dry tracts of Uttar Pradesh, Madhya Pradesh, Rajasthan, Punjab plains and Gujarat (Tripathi et al., 1998). This plant has ploidy level variations and predominance with tetraploid, chromosome number is $2n = 48$ (Iqbal and Dutta, 2007). It is a small erect, evergreen shrub, grown about 30-75 cm in height and covered with wooly pubescence. Roots are tuberous, 20-30 cm long, cylindrical, 1-2 cm diameter and whitish brown colour skin. The leaves are simple, opposite, ovate, and

glabrous, 4-10 cm length and 2-7 cm broad and arranged in alternate fashion. Flowers are bisexual, axillary umbellate cymes, five petals, and five sepals, bicarpellary ovary, single style, and bilobed stigma with dehiscent anthers (Atal and Schwarting, 1961; Kaul et al., 2005). The pollen load is huge on the stigma, and stiff pollen which favours self-pollination (Mir et al., 2012). Petals are united, stamens attached with corolla tube. The matured fruit is berry, orange-red in colour and is enclosed with persistent calyx. Seeds are tiny, reniform shape and yellow in colour (Bharti et al., 2016). Typical photographs of various botanical features of ashwagandha are presented in Figure 1.

Ethnopharmacological potential of ashwagandha

Ashwagandha roots have been used over 3000 years in traditional medicine for treating various health problems in Asia (Singh et al., 2010). This plant roots serves as ingredient for about 200 formulations in India folk medicine and is used as an aphrodisiac and for the treatment of asthma, inflammation, insomnia, anxiety, psoriasis, constipation, fatigue, weakness, impotence, premature ageing, ulcers, and mental stress. Ashwagandha leaves used for the treatment of tumors in Ayurvedic and Unani system of medicine (Singh et al., 2010). The plant root extracts and active molecule withanolides have potent therapeutic properties like antioxidant, immunomodulation, anti-ageing, adaptogenic, neuroprotective, and antitumor activities (Ziauddin et al., 1996; Rasool and Varalakshmi, 2006; Gupta et al., 2011; Dhuley, 2001; Mishra et al., 2000; Prakash et al., 2013 & 2014).

In Indian traditional medicine doctors and healers believed that ashwagandha root powder boiled with milk is recommended to cure sterility for females (Kirtikar and Basu, 1999). Traditionally, this plant roots used to stimulate youthful vigour and strength, improve the vital fluid production, blood muscle fat, semen and cells (Williamson, 2002). Ashwagandha fruits are used as bitter tonic for dyspepsia and promoting infants growth and possess potent agent for sedative, blood purified and diuretic (Watt, 1972). According to Patwardhan et al. (1998), ashwagandha leaves are recommended to cure the hurting swelling, fever and ophthalmitis. Furthermore, this plant also exposed antiserotogenic and anabolic properties and is displayed beneficial activities for the treatment of geriatrics, stress, and arthritis (Mirjalili et al., 2009; Alam et al., 2011). Also, in Indian folk medicine ashwagandha roots have been used for relaxing the mind and enhance the memory power and eye sight (John, 2014).



Figure 1. Botanical features of ashwagandha (*Withania somnifera* L.)




Figure 2. Ethnobotanical uses of *Withania somnifera*

Ashwagandha Rasayana is an extensively admired Ayurveda drug for peoples from the all ages. Consistent consumption of Ashwagandha is assumed to delay the senescence, correct the irregularities of sense organs, increase the growth of muscle cells, revitalize the reproductive organs and enhance the fertility. Furthermore, Ashwagandha rasayana is also useful to cure the cough, insomnia, muscular weakness, stress, memory loss, tiredness and aging problems (Sharma and Dash, 1998). Graphical representation of ethnobotanical uses of *Withania somnifera* has been shown in Figure 2.

CONCLUSIONS

Withania somnifera (L.) is an important medicinal plant and it has been used from ancient times in traditional Indian medicines (TIM). In Indian Ayurveda, this plant has grouped as Rasayana

(which means lengthening lifespan). All the *W. somnifera* plant parts has been used to cure several diseases by various countries particularly in India. Among the various plant parts, roots have been widely used for treatment of some diseases like tiredness, cough, insomnia, muscular strength, stress, memory loss, aging complications, arthritis, asthma, anxiety, constipation, ulcers sterility, and impotence. Several clinical studies need to confirm the biological activity of root powder, extracts and active chemical constituents for treatment of various diseases using animal and human models.

AUTHOR CONTRIBUTIONS

A.P & K.A wrote the manuscript. S.S., K.S.V., M.K & L.H collected data and P.S editing the manuscript.

No potential conflict of interest was reported by authors.

DISCLOSURE STATEMENT

Table 1. Role of *Withania somnifera* extracts of the different parts of the plant

Parts used	Type of extract	Types of activity	Diseases	Important Findings	References
Root	Methanol	Anticancer	Colan cancer	Increased ROS by chemotherapy	Henley et al. (2017)
Root	Alcohol	Anticancer	Murine B16F1 melanoma	Apoptosis by intrinsic pathway suppression	Sudeep et al. (2018)
Entire plants	Methanol (75%)	Stem cells	bone marrow cellularity increased	Increased total WBC	Kuttan (1996)
Root	Aqueous	Anticancer	Ascites carcinoma	7.5 Gy gamma radiation	Devi et al. (1996)
Root	Water	Anticancer	Human malignant melanoma cells	Nuclear blebbing	Halder et al. (2015)
Leaves	Methanol	Antiinflammatory	Inflammatory disorders	NF-kappa inhibition by preventing TNF-induced activation	Kaileh et al. (2007)
Leaves	Methanol	Breast, liver cancer	cancer cell lines	At S phase cell cycle arrested	Alfaifi et al. (2016)
Leaves	Water	Neuroinflammation	Inactivation of microglial	cell cycle arrest at G0/G1 and G2/M phase	Gupta and Kaur (2016)
Stem	Ethanol	Anticancer	HeLa, A549, BT474, MDA-MB-231	Multidrug Resistance Targeting	Maliyakkal et al. (2015)
Fruits	Methanol	Neurological	Disorders in related to Neurology	BBB (blood-brain-barrier) permeability	Vareed et al. (2014)
Compound	withaferian A	Antimicrobial	<i>Pseudomonas aeruginos</i> , <i>Staphylococcus aureus</i> , <i>Bacillus subtilus</i> , <i>Micrococcus pyogenes</i>	It showed strong activity against tested organisms	Abou-Douh, (2002)
Root	Water, Methanol	Antioxidant	Chemo-preventive activity	Free radical scavenging action of the extract	Prakash et al. (2002)

DISCLOSURE STATEMENT

No potential conflict of interest was reported by authors.

REFERENCES

Abou-Douh, A.M. (2002). New withanolides and other constituents from the fruit of *Withania somnifera*. *Arch Pharm (Weinheim)*. 335, 267-276.

Alam, N., Hossain, M., Mottalib, Md A., Sulaiman, S.A., Gan, S.H. & Khalil, Md. I. (2012). Methanolic extracts of *Withania somnifera* leaves, fruits and roots possess antioxidant properties and antibacterial activities. *BMC Complementary and Alternative Medicine*, 12, 175. doi: 10.1186/1472-6882-12-175.

Alfaifi, M.Y., Saleh, K.A., El-Boushnak, M.A., Elbehairi, S.E., Alshehri, M.A. & Shati, A.A. (2016). Antiproliferative activity of the methanolic extract of *withania somnifera* leaves from faifa

- mountains, southwest saudi arabia, against several human cancer cell lines. *Asian Pacific Journal of Cancer Prevention*, 17, 2723–2726.
- Atal, C.K. & Schwarting, A.E. (1961). Aswagandha, an ancient Indian drug. *Economic Botany*, 15(3), 256–263.
- Withania somnifera*. *International Journal of Radiation Biology*, 69, 193–197.
- Dhuley, J.N., (2001). Nootropic-like effect of ashwagandha (*Withania somnifera* L.) in mice. *Phytotherapy Research*, 15(6), 524–528.
- Durg, S., Shivaram, S.B. & Bavage, S. (2018). *Withania somnifera* (Indian ginseng) in male infertility: An evidence-based systematic review and meta-analysis. *Phytomedicine*, 50, 247–256.
- Gupta, M., Bisht, D., Pandey, M.M., Ojha, S.K., Khatoon, S., Rastogi, S. & Rawat, A.K.S. (2011). Standardization of Ashwagandhilehya - An important ayurvedic formulation of *Withania somnifera*. *India Journal of Traditional Knowledge*, 10, 594–598.
- Gupta, M. & Kaur, G. (2016). Aqueous extract from the *Withania somnifera* leaves as a potential anti-neuroinflammatory agent: A mechanistic study. *Journal of Neuroinflammation*, 13(1), 193. doi: 10.1186/s12974-016-0650-3.
- Halder, B., Singh, S. & Thakur, S.S. (2015). *Withania somnifera* root extract has potent cytotoxic effect against human malignant melanoma cells. *PLoS ONE*, 10, e0137498.
- Henley, A.B., Yang, L., Chuang, K.L., Sahuri-Arisoylu, M., Wu, L.H., Bligh, S.W. & Bell, J.D. (2017). *Withania somnifera* root extract enhances chemotherapy through ‘priming’. *PLoS ONE*, 12, e0170917.
- Iqbal, M. & Dutta, A.M. (2007). Cytogenetic Studies in *Withania somnifera* (L.) Dun. (Solanaceae). *Cytologia*, 72(1), 43–47.
- John, J. (2014). Therapeutic potential of *Withania somnifera*: a report on phytopharmacological properties. *International Journal of Pharmaceutical Sciences and Research*, 5(6), 2131–2148.
- Kaul, M.K., Kumar, A. & Sharma A. (2005). Reproductive biology of *Withania somnifera* (L.) Dunal. *Current Science*, 88(9), 1375–1377.
- Kirtikar, K.R. & Basu, B.D. (1999). Indian Medicinal Plants. Vol.3, International Book Distributors Book Sellers and Publishers, Deheradun.
- Kulkarni, S.K. & Dhir A. (2008). *Withania somnifera*: an Indian ginseng. *Progress in Neuropsychopharmacology and Biological Psychiatry*, 32, 1093–105.
- Bharti V.K., Malik, J.K. & Gupta, R.C. (2016). Ashwagandha: multiple health benefits. *Nutraceuticals*, 717–733.
- Devi, P.U., Akagi, K., Ostapenko, V., Tanaka, Y., Sugahara, T. (1996). Withaferin A: A new radio sensitizer from the Indian medicinal plant
- Kuttan, G. (1996). Use of *Withania somnifera* Dunal as an adjuvant during radiation therapy. *Indian Journal of Experimental Biology*, 34, 854–856.
- Maliyakkal, N., Appadath Beeran, A., Balaji, S.A., Udupa, N., Ranganath Pai, S. & Rangarajan, A. (2015). Effects of *Withania somnifera* and *Tinospora cordifolia* extracts on the side population phenotype of human epithelial cancer cells: Toward targeting multidrug resistance in cancer. *Integrative Cancer Therapies*, 14, 156–171.
- Mir, B.A., Koul, S., Kumar, A., Sushant, S., Kaul, M.K. & Soodan A.S. (2012). Reproductive behaviour and breeding system of wild and cultivated types of *Withania somnifera* (L.) Dunal. *Journal of Medicinal Plants Research*, 6(5), 754–762.
- Mirjalili, M.H., Moyano, E., Bonfill, M., Cusido, R.M. & Palazon, J. (2009). Steroidal lactones from *Withania somnifera*, an ancient plant for novel medicine. *Molecules*, 14(7), 2373–2393.
- Mishra, LC., Singh, B.B. & Dagenais, S. (2000). Scientific basis for the therapeutic use of *Withania somnifera* (Ashwagandha): A review. *Alternative Medicine Review*, 5, 334 – 346.
- Patwardhan, B., Panse, G.T. & Kulkarni, P.H. (1998). Ashwagandha a review. *Journal of National Integrative Medical Association*, 30, 7–11.
- Prakash, J., Chouhan, S., Yadav, S.K., Westfall, S., Rai, S.N. & Singh, S.P. (2014). *Withania somnifera* alleviates parkinsonian phenotypes by inhibiting apoptotic pathways in dopaminergic neurons. *Neurochemical Research*, 12, 2527–2536.
- Prakash, J., Gupta, S.K. & Dinda, A.K. (2002). *Withania somnifera* root extract prevents dmba-induced squamous cell carcinoma of skin in Swiss albino mice. *Nutrition and Cancer*, 42, 91–97.
- Prakash, J., Yadav, S.K., Chouhan, S. & Singh, S.P. (2013). Neuroprotective role of *Withania somnifera* root extract in maneb-paraquat induced mouse model of parkinsonism. *Neurochemical Research*, 38(5), 972–80.
- Rajasankar, S., Manivasagam, T., Sankar, V., Prakash, S., Muthusamy, R., Krishnamurti, A. & Surendran, S. (2009). *Withania somnifera* root extract improves catecholamines and physiological abnormalities seen in a

- Parkinson's disease model mouse. *Journal of Ethnopharmacology*, 125, 369-373.
- Rasool, M. & Varalakshmi, P. (2006). Immunomodulatory role of *Withania somnifera* root powder on experimental induced inflammation: An *in vivo* and *in vitro* study. *Vascular Pharmacology*, 44, 406 - 410.
- Sharma, R.K. & Dash, B. (1998). *Carka Samhita*, Vol. II. Varanasi, India: Chowkamba Sanskrit Series Office.
- Singh, G., Sharma, P.K., Dudhe, R. & Singh, S. (2010). Biological activities of *Withania somnifera*. *Annals of Biological Research*, 1(3), 56-63.
- Singh, R.H., Narsimhamurthy, K. & Singh, G. (2008). Neuronutrient impact of Ayurvedic Rasayana therapy in brain aging. *Biogerontology*, 9, 369-374.
- Sudeep, H.V., Gouthamchandra, K., Venkatesh, B.J., Prasad, K. & Viwithan, S (2018). A standardized *Withania somnifera* root extract induces apoptosis in murine melanoma cells. *Pharmacognosy Magazine*, 13, S801-S806.
- Tripathi, A.K., Dey, S., Singh, R.H. & Dey, P.K. (1998). Alterations in the sensitivity of 5th receptor subtypes following chronic Ashwagandha treatment in rats. *Ancient Sciences of Life*, 17(3):168-181.
- Vareed, S.K., Bauer, A.K., Nair, K.M., Liu, Y., Jayaprakasam, B. & Nair, M.G. (2014). Blood-brain barrier permeability of bioactive withanamides present in *Withania somnifera* fruit extract. *Phytotherapy Research*, 28, 1260-1264.
- Watt, G.A. (1972). *Dictionary of the economic Products of India*. Cosmo Publication, Delhi, India.
- Williamson, E.M. (2002). *Major herbs of Ayurveda*, Churchill Livingstone: London, UK, p.322-323.
- Ziauddin, M., Phansalkar, N., Patki, P., Diwanay, S. & Patwardhan, B. (1996). Studies on the immunomodulatory effects of Ashwagandha. *Journal of Ethnopharmacology*, 50, 69 - 76.